


Generation of digital planes using generalized continued-fractions algorithms

Xavier Provençal

Joint work with : Damien Jamet and Nadia Lafrenière

Journées nationales du GDR IM

Jeudi le 16 mars, Montpellier



Arithmetic digital line

Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms



Definition (Reveillès (1991), Kovalev (1990))

An **arithmetic digital line** is the set :

$$\mathcal{D}((a, b), \mu) = \{(x, y) \in \mathbb{Z}^2 \mid 0 \leq ax + by + \mu < |a| + |b|\}$$

where

- (a, b) is the **normal vector**,
- $-b/a$ is the **slope**,
- μ is the **shift**.



Digital Straight Segment (DSS)

Periodic
structure


Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

Definition


A **digital straight segment** is a finite and connected subset of a digital line.



Tangential cover

Definition ([Feschet, Tougne 99])

The **tangential cover** of a discrete shape is the sequence of all maximal DSS on its boundary.



Theorem ([Debled-Rennesson, Reveilles 1995][Lachaud, vialard, de Vieilleville 2007])

The computation of the tangential cover take a time in $\mathcal{O}(n)$ where n is the number of points on the boundary of the shape.

Applications of the tangential cover include :

- Convexity test
[Debled-Rennesson, Reiter-Doerksen 04]
- Tangent estimation
[Feschet, Tougne 99],
[Lachaud, de Vieilleville 07]
- Length estimation
[Lachaud, de Vieilleville 07]
- Curvature estimation
[Lachaud, Kerautret, Naegel 08]
- Automatic noise detection
[Lachaud, Kerautret 12]

Digital lines and planes

Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms




Definition ([Reveillès 91])

The **digital line/plane/hyperplane** $\mathcal{P}(v, \mu, \omega)$ with **normal vector** $v \in \mathbb{Z}^d$, **thickness** $\omega \in \mathbb{N}$ and **shift** $\mu \in \mathbb{R}$ is the subset of \mathbb{Z}^d defined by:

$$\mathcal{P}(v, \mu, \omega) = \{x \in \mathbb{Z}^d \mid 0 \leq \langle x, v \rangle - \mu < \omega\}$$

$$\mathcal{P}((1, 6), 7, 0)$$

$$0 \leq 1x + 6y < 7$$



Digital lines and planes

Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms




Definition ([Reveillès 91])

The **digital line/plane/hyperplane** $\mathcal{P}(v, \mu, \omega)$ with **normal vector** $v \in \mathbb{Z}^d$, **thickness** $\omega \in \mathbb{N}$ and **shift** $\mu \in \mathbb{R}$ is the subset of \mathbb{Z}^d defined by:

$$\mathcal{P}(v, \omega) = \{x \in \mathbb{Z}^d \mid 0 \leq \langle x, v \rangle < \omega\}$$

$$\mathcal{P}((1, 6), 7)$$


$$0 \leq 1x + 6y < 7$$



Periodic structure of a digital line

Example with $v = (-3, 1)$:


- $\langle x, v \rangle$ is the **height** of x ,
- $\mathcal{P}(v, 4) = \{x \in \mathbb{Z}^2 \mid 0 \leq \langle x, v \rangle < 4\}$.



Periodic structure of a digital line

Example with $v = (-3, 1)$:



- $\langle x, v \rangle$ is the **height** of x ,
- $\mathcal{P}(v, 4) = \{x \in \mathbb{Z}^2 \mid 0 \leq \langle x, v \rangle < 4\}$.



- $\langle x, v \rangle = \langle y, v \rangle \implies y - x$ is a period vector.
- A point of each height from 0 to $\|v\|_1 - 1$ appear in a period.

Periodic structure of a digital plane

$$v = (1, 2, 3), \quad \mathcal{P}(v, 6) = \{x \in \mathbb{Z}^3 \mid 0 \leq \langle x, v \rangle < 6\}$$



Periodic
structure



Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

Periodic structure of a digital plane

$$v = (1, 2, 3), \quad \mathcal{P}(v, 6) = \{x \in \mathbb{Z}^3 \mid 0 \leq \langle x, v \rangle < 6\}$$



Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

Periodic structure of a digital plane



Periodic structure

Construction guided by Euclid

Fully Subtractive

New algorithms

$$v = (1, 2, 3), \quad \mathcal{P}(v, 6) = \{x \in \mathbb{Z}^3 \mid 0 \leq \langle x, v \rangle < 6\}$$



Periodic structure of a digital plane



Periodic structure

Construction guided by Euclid

Fully Subtractive

New algorithms

$$v = (1, 2, 3), \quad \mathcal{P}(v, 6) = \{x \in \mathbb{Z}^3 \mid 0 \leq \langle x, v \rangle < 6\}$$



- $\langle x, v \rangle = \langle y, v \rangle \implies y - x$ is a period vector.

Periodic structure of a digital line

Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms


Definition

A set of points $S \subset \mathbb{Z}^d$ provided with a set of vectors $(b_i)_{i=1}^n \in \mathbb{Z}^d$ covers an infinite set $\Omega \subset \mathbb{Z}^d$ if

$$\Omega = \bigcup_{x \in \mathbb{Z}b_1 + \mathbb{Z}b_2 + \dots + \mathbb{Z}b_n} (S + x).$$

(Like a tiling without a disjoint union.)

Example :



provided with vector $v = (3, 1)$ covers the digital line $\mathcal{P}((-3, 1), 4)$.

Main pattern of a digital line


Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

- A point $x \in \mathcal{P}(v, \|v\|_1)$ is a **upper leaning point**, noted **UL**, if its height $\langle x, v \rangle$ is maximal.



Main pattern of a digital line


Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

- A point $x \in \mathcal{P}(v, \|v\|_1)$ is a **upper leaning point**, noted **UL**, if its height $\langle x, v \rangle$ is maximal.
- The **main pattern** of a digital line is a set of points bounded by two consecutive upper leaning points.



Main pattern of a digital line


Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms


- A point $x \in \mathcal{P}(v, \|v\|_1)$ is a **upper leaning point**, noted **UL**, if its height $\langle x, v \rangle$ is maximal.
- The **main pattern** of a digital line is a set of points bounded by two consecutive upper leaning points.
- Let v be the vector defined by two consecutive **UL**, a main pattern provided with v covers its digital line.



Main pattern of a digital line

- \circ : upper leaning points.
- Let H be the highest point among $\{\bullet\}$.


Main pattern of slope $2/5$.



Main pattern of a digital line

- \circ : upper leaning points.
- Let H be the highest point among $\{\bullet\}$.


Main pattern of slope $2/5$.



Main pattern of a digital line

- \circ : upper leaning points.
- Let H be the highest point among $\{\bullet\}$.



Main pattern of slope $2/5$.



Main pattern of a digital line

- \circ : upper leaning points.
- Let H be the highest point among $\{\bullet\}$.

Main pattern of slope $2/5$.





Main pattern of slope $3/8$.


Main pattern of a digital line

- \circ : upper leaning points.
- Let H be the highest point among $\{\bullet\}$.

Main pattern of slope $2/5$.



Main pattern of slope $3/7$.



Main pattern of slope $3/8$.


Stern-Brocot Tree

Periodic structure


Construction guided by Euclid

Fully Subtractive

New algorithms




Stern-Brocot tree.



Every irreducible fraction appears exactly once in the Stern-Brocot tree.

Stern-Brocot tree



Euclid Algorithm

Euclid algorithm

Approximation

(7, 9)

(1, 1)

↓

↓

(7, 2)

(1, 2)

↓

↓

(5, 2)

(2, 3)

↓

↓

(3, 2)

(3, 4)

↓

↓

(1, 2)

(4, 5)

↓

↓

(1, 1)

(7, 9)

Matricial view

Periodic structure

Construction guided by Euclid

Fully Subtractive

New algorithms

	Euclid algorithm	Approx.
n	v_n	a_n
0	(<u>7</u> , 9)	(1, 1)
	↓	↓
1	(7, <u>2</u>)	(1, 2)
	↓	↓
2	(5, <u>2</u>)	(2, 3)
	↓	↓
3	(3, <u>2</u>)	(3, 4)
	↓	↓
4	(<u>1</u> , 2)	(4, 5)
	↓	↓
5	(1, 1)	(7, 9)

Euclid algorithm

Given a vector (x, y) , return

- $\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix}$ if $x < y$,
- $\begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}$ if $x > y$,
- **stop** if $x = y$.

Given a vector $v \in (\mathbb{N} \setminus \{0\})^2$, let :

- $v_0 = v$,
- For all $n \geq 1$: $\begin{cases} M_n = \mathbf{Euclid}(v_{n-1}) \\ v_n = M_n v_{n-1}. \end{cases}$

Property


- $v_n = M_n M_{n-1} \cdots M_1 v$
- $a_n = M_1^{-1} M_2^{-1} \cdots M_n^{-1} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$

Matricial view

Let UL_0 and UL_1 be two upper leaning points of a main pattern of $\mathcal{P}(a_n, \|a_n\|_1)$ and H be the Bezout point. Let $\alpha = UL_0 - H$ and $\beta = UL_1 - H$, then

$$M_1^T M_2^T \cdots M_n^T = \begin{bmatrix} \alpha & \beta \end{bmatrix}$$

$$M_1^T \cdots M_n^T e_1 = \alpha, \quad M_1^T \cdots M_n^T e_2 = \beta.$$



The Translation-Union Construction

Periodic structure

Construction guided by Euclid

Fully Subtractive

New algorithms

Construction

[Domenjoud, Vuillon 12],
[Berthé, Jamet, Jolivet, P. 2013]

Let $v_0 = v$, $B_0 = \{\mathbf{0}\}$ and for all $n \geq 1$ let :

M_n : the matrix selected from v_{n-1} ,

$$v_n = M_n v_{n-1}$$

δ_n : the index of the coordinate of v_{n-1} that is subtracted,

$$T_n = M_1^T \cdots M_n^T e_{\delta_n}, \quad (\text{translation})$$

$$B_n = B_{n-1} \cup (T_n + B_{n-1}), \quad (\text{body})$$

$$H_n = \sum_{i \in \{1, \dots, n\}} T_i, \quad (\text{highest point})$$

$$L_n = H_n + \{M_1^T \cdots M_n^T e_i\}. \quad (\text{legs})$$


Note that:

$$H_n \in B_n,$$


$$L_n \cap B_n = \emptyset.$$

$$\bullet \in B_n, \quad \circ \in L_n$$


$$\begin{aligned} v_0 &= (2, 3), \\ a_0 &= (1, 1) \\ H_0 &= (0, 0), \\ L_0 &= \{(1, 0), (0, 1)\}. \end{aligned}$$



$$\begin{aligned} v_1 &= (2, 1), \delta_1 = 1 \\ a_1 &= (1, 2) \\ T_1 &= (1, 0) \\ H_1 &= (1, 0), \\ L_1 &= \{(2, 0), (0, 1)\}. \end{aligned}$$



$$\begin{aligned} v_2 &= (1, 1), \delta_2 = 2 \\ a_2 &= (2, 3) \\ T_2 &= (-1, 1) \\ H_2 &= (0, 1), \\ L_2 &= \{(2, -1), (-1, 1)\}. \end{aligned}$$



3D continued fraction algorithms

Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

Euclid algorithm : given two numbers subtract the smallest to the largest.

$$(7, 9) \rightarrow (7, 2) \rightarrow (5, 2) \rightarrow (3, 2) \rightarrow (1, 2) \rightarrow (1, 1) \rightarrow (1, 0) \curvearrowright$$

Given three numbers :

- **Selmer** : subtract the smallest to the largest.
 $(3, 7, 5) \rightarrow (3, 4, 5) \rightarrow (3, 4, 2) \rightarrow (3, 2, 2) \rightarrow (1, 2, 2) \rightarrow (1, 2, 0) \curvearrowright .$
- **Brun** : subtract the second largest to the largest.
 $(3, 7, 5) \rightarrow (3, 2, 5) \rightarrow (3, 2, 2) \rightarrow (1, 2, 2) \rightarrow (1, 2, 0) \rightarrow (1, 1, 0) \rightarrow (1, 0, 0) \curvearrowright .$
- **Fully subtractive** : subtract the smallest to the two others.
 $(3, 7, 5) \rightarrow (3, 4, 2) \rightarrow (1, 2, 2) \rightarrow (1, 1, 1) \rightarrow (1, 0, 0) \curvearrowright .$
- **Poincaré** : subtract the smallest to the mid and the mid to the largest.
 $(3, 7, 5) \rightarrow (3, 2, 2) \rightarrow (1, 2, 0) \rightarrow (1, 1, 0) \rightarrow (1, 0, 0) \curvearrowright .$
- **Arnoux-Rauzy** : subtract the sum of the two smallest to the largest (not always possible).
 $(3, 7, 5) \rightarrow$ impossible.
- ...

Example : Fully Subtractive $v = (6, 8, 11)$

Construction

Let $v_0 = v$, $B_0 = \{0\}$ and for all $n \geq 1$ let :

M_n : the matrix selected from v_{n-1} ,

$$v_n = M_n v_{n-1}$$

δ_n : the index of the coordinate of v_{n-1} that is subtracted,


$$T_n = M_1^T \cdots M_n^T e_{\delta_n}, \quad (\textit{translation})$$

$$B_n = B_{n-1} \cup (T_n + B_{n-1}), \quad (\textit{body})$$


$$H_n = \sum_{i \in \{1, \dots, n\}} T_i, \quad (\textit{highest point})$$

$$L_n = H_n + \{M_1^T \cdots M_n^T e_i\}. \quad (\textit{legs})$$


- Step 0 : $v_0 = (6, 8, 11)$, $a_0 = (1, 1, 1)$,



- Step 1 : $v_1 = (6, 2, 5)$, $a_1 = (1, 2, 2)$,



- Step 2 : $v_2 = (4, 2, 3)$, $a_2 = (2, 3, 4)$,



Example : Fully Subtractive $v = (6, 8, 11)$


Periodic
structure

Construction
guided by
Euclid


Fully
Subtractive


New
algorithms

- Step 3 : $v_3 = (2, 2, 1)$, $a_3 = (3, 4, 6)$,



- Step 4 : $v_4 = (1, 1, 1)$, $a_4 = (6, 8, 11)$,






$\mathcal{P}((6, 8, 11), 13)$

Expected properties of the pattern:

- Connected.
- Provides period vectors.
- Covers $\mathcal{P}(v, \omega)$ with these vectors.
- Should be as small as possible, to avoid redundancy.



Example, Fully Subtractive $v = (6, 8, 13)$

Periodic structure

Construction guided by Euclid


Fully Subtractive

New algorithms


- Step 0 : $v_0 = (6, 8, 13)$, $a_0 = (1, 1, 1)$,




- Step 1 : $v_1 = (6, 2, 7)$, $a_1 = (1, 2, 2)$,




- Step 2 : $v_2 = (4, 2, 5)$, $a_2 = (2, 3, 4)$,



- Step 3 : $v_3 = (2, 2, 3)$, $a_3 = (3, 4, 6)$,



- Step 4 : $v_4 = (2, 0, 1)$, $a_4 = (5, 7, 11)$,



Definition

Let \mathcal{K} be the set of vectors v such $\mathbf{FS}^N(v) = (1, 1, 1)$ for some $N \geq 1$.

Let $v \in (\mathbb{N} \setminus \{0\})^3$ with $\gcd(v) = 1$ and $(a, b, c) = \text{sort}(v)$ (i.e. $a \leq b \leq c$), two conditions:

- (1) If $a + b \leq c$ then let $(a', b', c') = \text{sort}(\mathbf{FS}(v))$ then $a' + b' \leq c'$.

$$\text{Example : } (2, 3, 6) \xrightarrow{\mathbf{FS}} (2, 1, 4) \xrightarrow{\mathbf{FS}} (1, 1, 3) \xrightarrow{\mathbf{FS}} (1, 0, 2).$$

- (2) If $a = b < c$, then one coordinate of $\mathbf{FS}(v)$ is 0.


$$\text{Example : } (2, 2, 3) \xrightarrow{\mathbf{FS}} (2, 0, 1).$$

Lemma

Let $v \in (\mathbb{N} \setminus \{0\})^3$, $v \notin \mathcal{K}$ iff there exist $n \geq 0$ such that $\mathbf{FS}^n(v)$ satisfies condition (1) or (2).

The set \mathcal{K}

$$v \xrightarrow{\text{FS}} \dots \xrightarrow{\text{FS}} (1, 1, 1)$$



Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

New generalized continued fraction algorithms

Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

Idea : If the vector *looks good*, use **FS**, otherwise use some thing else. . . like **Brun** or **Selmer**.

New generalized continued fraction algorithms

Periodic
structure

Construction
guided by
Euclid

Fully
Subtractive

New
algorithms

Idea : If the vector *looks good*, use **FS**, otherwise use some thing else... like **Brun** or **Selmer**.

Algorithm FSB
Input : $v \in \mathbb{N}^3$.
If v satisfies (1) or (2) then Use Brun . else Use FS . end if

Algorithm FSS
Input : $v \in \mathbb{N}^3$.
If v satisfies (1) or (2) then Use Selmer . else Use FS . end if

Example using **FSB**, $v = (9, 15, 11) \notin \mathcal{K}$

Periodic structure

Construction guided by Euclid

Fully Subtractive

New algorithms

$$v_0 = (9, 15, 11)$$


$$a_0 = (1, 1, 1)$$



FS →

$$v_1 = (9, 6, 2)$$

$$a_1 = (1, 2, 2)$$



Brun →

$$v_2 = (3, 6, 2)$$

$$a_2 = (2, 3, 3)$$



Brun →

$$v_3 = (3, 3, 2)$$


$$a_3 = (3, 5, 4)$$



FS →

$$v_4 = (1, 1, 2)$$

$$a_4 = (6, 10, 7)$$



Theorem


Using the algorithm **FSB** or **FSS**, for all vector $v \in (\mathbb{N} \setminus \{0\})^3$ with $\gcd(v) = 1$,

- ① $\exists N$ such that $v_N = (1, 1, 1)$.
- ② Vectors of L_N have same height, providing period vectors.
- ③ $B_N \cup L_N$ is connected.
- ④ $B_N \cup L_N$ covers $\mathcal{P}(v, \omega)$ with $\frac{\|v\|_1}{2} \leq \omega < \|v\|_1$.

Brun \rightarrow

$$v_5 = (1, 1, 1)$$

$$a_5 = (9, 15, 11)$$




Conclusion

Periodic structure

Construction guided by Euclid

Fully Subtractive


New algorithms



Good:

- Build a pattern that covers a digital plane for any rational normal vector.
- Construction is recursive and based on continued fractions algorithms.
- Generalizes Voss' *splitting formula* (equiv. *standard factorization* of Christoffel words) to higher dimensions.

$\mathcal{P}((9, 15, 11), 23)$



Problems: Open questions :

- Find a gcd algorithm that builds minimal patterns.
- Control the height of the pattern.
- Control the anisotropy of the patterns (avoid stretched forms in favor of *potato-likeness*).
- Apply recursive structure to image analysis algorithms.